

# Conservation of non-canonical start sites in FMR-1

Dionne Weaver, Gianni Walker, and Dr. Jack Vincent

## Introduction:

While working on a previous annotation project, evidence for a non-canonical (or non-AUG) start site was found in two isoforms for the *FMR-1* gene in the fruit fly species, *Drosophila erecta*. Upon further research, we found a paper that discussed the conservation of the 5' UTR within the *melanogaster* subgroup. We then wondered if this conservation also meant the non-canonical start site was conserved and if so, how far into the *Drosophila* genus it went.

This project looked for evidence a non-canonical start site within the same isoforms of four additional species. The species looked at were *D. ficusphila*, *D. obscura*, *D. arizonae*, and *D. buskii* and were chosen based on their evolutionary split from *D. melanogaster* whose genome has been annotated.

		Second letter					
		U	C	A	G		
U	UUU	Phe	UCU	UAC	Tyr	UGU	Cys
	UUC		UCC	UAG	Stop	UGC	Stop
	UUA	Leu	UCA	UAA	Stop	UGA	Stop
	UUG		UCG	UAG	Stop	UGG	Trp
C	CUU		CCU	CAU	His	CGU	Arg
	CUC		CCC	CAC	CGC		
	CUA	Leu	CCA	CAA	Gln	CGA	
	CUG		CCG	CAG		CGG	
A	AUU		ACU	AAU	Asn	AGU	Ser
	AUC	Ile	ACC	AAC	AGC	AGA	
	AUA		ACA	AAA	Lys	AGG	
	AUG	Met	ACG	AAG		AGG	
G	GUU		GCU	GAU	Asp	GGU	Gly
	GUC		GCC	GAC		GGC	
	GUA	Val	GCA	GAA		GGA	
	GUG		GCG	GAG		GGG	

Green: typical AUG start codon

Red: non-canonical start codon found within the G and H isoform of *D. melanogaster*

For more information please contact:

DionneW@UW.EDU, GWalker9@UW.EDU, or JaV2@UW.EDU

## Objectives:

Our objectives for this project were to 1. annotate the *FMR-1* gene in the four chosen species and 2. analyze the G and H isoforms for evidence of the non-canonical start site.


## Materials & Methods:

The GEP UCSC genome browser was first used to examine the genomic neighborhood of *D. melanogaster*. The predicted protein sequence for the *FMR-1* gene was used as the query for the tblastn search to find the ortholog in the four chosen species. Once the ortholog was found, the genomic neighborhood was examined. Using the accession numbers from the refseq BLAST alignment we obtained the predicted protein sequences and compared them to that of *D. melanogaster*. After visual confirmation of the ortholog, the gene record finder was used to find the approximate locations of the coding exons (CDS's). The location of the CDS's was then refined using a BLAST search comparing the sequence within *D.*

*melanogaster* to the target species. Using the genome browser, we then visually examined each exon for evidence of the non-canonical start sites, focusing on the G and H isoforms. We looked for any strong indications of homology between the BLAT, SPALN, and refseq data. The potential non-canonical start site was confirmed by both the CUG codon as well as support from RNAseq data.

## Results:

Of the four species examined, only one species, *D. ficusphila* showed evidence of the non-canonical start site.



## References:

Beerman RW, Jongens TA. 2011. A non-canonical start codon in the *Drosophila* fragile X gene yields two functional isoforms. *Neuroscience*. 181:48–66.

## Acknowledgements:

Genome Pathways Project for giving us a research topic  
University of Washington Tacoma Sciences department for the opportunity to do a research project